

3.2.1 Number of papers published per teacher in the Journals notified on UGC (UGC CARE LIST JOURNALS) website during the year 2021-22

Sr. no	Title of paper	Name of the author/s	Department of the teacher	Name of journal	Year of publication	ISSN number	Link to the recognition in UGC enlistment of the Journal
1	Seasonal Fluctuation of Rice Brown Planthopper (Homoptera: Delphacidae), <i>Nilaparvata lugens</i> (Stal) - A Major Paddy Pest in Central India	Madhukar F. Jadhao and Arti A. Salwe	Zoology	Journal of Scientific Research	2021-22	0447-9483	<p>Journal Link- https://internet.bhu.ac.in/research_pub/jsr/Volume_65_06_2021.html</p> <p>Paper Link- https://internet.bhu.ac.in/research_pub/jsr/Volumes/JSR_65_06_2021/Manuscript%2025.pdf</p>
2	Biology of a Pentatomid Bug <i>Amyotea malabarica</i> (Fabricius) and its Predatory Efficiency in Rice Field of Vidarbha, India	Barsagade D.D. and Salwe A. A.	Zoology	International Journal of Zoological Investigations	2021-22	2454-3055	<p>Journal Link- https://mjl.clarivate.com/:search-results?issn=2454-3055&hide_exact_match_fl=true&utm_source=mjl&utm_medium=share-by-link&utm_campaign=search-results-share-this-journal</p> <p>Paper Link- 4514553453IssueSpecial 4 (1).pdf</p>
3	Sahitya Vishayak Krantikari Bhumikechi Awashyakata	Renukadas Ubale	Marathi	Parivartanacha Watsaru	2021-22	2250-3145	http://www.pvatsaru.com/pvatsaruweb/frontend/article/show/3570



Seasonal Fluctuation of Rice Brown Planthopper (Homoptera: Delphacidae), *Nilaparvata lugens* (Stal) - A Major Paddy Pest in Central India

Madhukar F. Jadhao^{*1} and Arti A. Salwe²

*¹ & ²Department of Zoology, S. N. Mor College, Tumsar- 441912, Dist- Bhandara (Maharashtra), India

mfjadhao@gmail.com, aaszoology@gmail.com

Abstract: Among the 5 species of sap feeders belonging to order Homoptera identified in the present investigation in both rabi and kharif seasons, *Nilaparvata lugens* was infesting rice crop very commonly and thus representing major pest status in Bhandara district (a rice bowl of state) of Maharashtra (central India). The overall population growth rate and peak density of *N. lugens* during rabi season was much lower as compared to the kharif season. The peak population of *N. lugens* was observed during the late kharif seasons from October to November. Another peak appeared during the rabi season from April to May. The infestation of *N. lugens* appeared in appreciable number at the grain filling stage of crop. The correlation analysis study showed that in rabi season, *N. lugens* exhibited highly positive correlation with relative humidity and significant negative correlation with maximum temperature whereas it showed highly significant negative correlation with relative humidity and minimum temperature during kharif season. The linear regression equations derived from the data may help in predicting the occurrence of these major pests in rice ecosystem of this region.

Keywords: Rabi, Central India, Correlation, Kharif, *Nilaparvata lugens*, Paddy pest, Regression.

I. INTRODUCTION

In India, rice (paddy), *Oryza sativa* L. is a staple food for over 55% population and is grown in almost all the states. Despite having the largest area under rice, its yield is among the lowest in the world (Rangi, 1993). The insect pests constitute one of the major yields limiting biotic stresses for rice crop throughout the world. About 300 species of insects have been reported to attack rice crop in India, of which 20 have been

found to be the major pests (Pathak, 1967) causing 21 to 51 percent yield losses (Singh and Dhaliwal, 1994). The present paper describes the seasonal incidence and correlation between weather parameters and population of a major pest, *Nilaparvata lugens* in the rice ecosystem of Bhandara district of eastern Maharashtra (central India) in relation to climatic conditions of this region.

extensively in the oil, gas and chemical industries because of its outstanding mechanical properties. The use of steel is also one of the effective strategies to maximize profit and reduce cost as compared to expensive corrosion resistant alloys. In many industrial applications related to oil and gas processing such as pipeline cleaning, pipeline/acid descaling and oil well acidizing, the use of mineral acids (usually hydrochloric acid) is still an effective method for improving productivity. This process however endangers the life of steel structures as a result of acid driven corrosion. In order to prevent this undesirable reaction, corrosion inhibitors are often added to the acid solution during acidification process (Hassan AR., and Gbadeyan JA., 2015; Ferdows M., et al.2009). Organic compounds, containing functional electronegative groups and π -electron in triple or conjugated double bonds are usually good inhibitors (Cano E., et al 2004; Ghailane T., et al.2013). Heteroatoms, such as sulphur, phosphorus, nitrogen and oxygen, together with aromatic rings in their structure are the major adsorption centres (Goulart C M., et al.2013; Bentiss F., et al 2005). The planarity (g) and the lonely electron pairs in the heteroatoms are important features that determine the adsorption of these molecules on the metallic surface (Hegazy MA., et al.2013; Hu SQ., et al.2010). The use of corrosion inhibitors is one of the most effective measures for

https://ugccare.unipune.ac.in/Ap... 80% ⋮

≡ UGC- Cell for Journal Analysis

Journal Details

Home **UGC** **Search**

Journal Title (in English Language)	Journal of Scientific Research
Publication Language	English
Publisher	Institute of Science, Banaras Hindu University
ISSN	0447-9483
E-ISSN	NA
Discipline	Science
Subject	Agricultural and Biological Sciences (all) , Biochemistry, Genetics and Molecular Biology (all) , Chemistry (all) , Engineering (all) , Mathematics (all) , Medicine (all)
Focus Subject	Agricultural and Biological Sciences (miscellaneous) , Biochemistry, Genetics and Molecular Biology (miscellaneous) , Chemistry (miscellaneous) , Engineering (miscellaneous) , Mathematics (miscellaneous) , Medicine (miscellaneous)
UGC-CARE List Group	Group D

Copyright © 2019 Savitribai Phule Pune University. All rights reserved. | Disclaimer

Volume 7 Special Issue 2021

ISSN 2454-3055

Manuscripts published in Special Issue were presented at National Virtual Conference on "Current Research Trends In Zoology" on 5-6 August, 2021 at Department of Zoology, Elphinstone College, Mumbai, India

INTERNATIONAL JOURNAL OF ZOOLOGICAL INVESTIGATIONS

*Forum for Biological and
Environmental Sciences*

Published by Saran Publications, India



Biology of a Pentatomid Bug *Amyotea malabarica* (Fabricius) and its Predatory Efficiency in Rice Field of Vidarbha, India

Barsagade D.D.¹ and Salwe A. A.^{2*}

¹P.G. Department of Zoology, M.J.F. Educational Campus, R.T.M. Nagpur University, Nagpur 440 033, India

²Department of Zoology, S. N. Mor College of Arts, Commerce and Smt. G. D. Saraf Science College, Tumsar 441 912 (M.S.), India

*Corresponding Author

Received: 14th August, 2021; Accepted: 30th August, 2021; Published online: 5th September, 2021

<https://doi.org/10.33745/ijzi.2021.v07i02s.004>

Abstract: *Amyotea malabarica* (Fabricius) (Heteroptera: Pentatomidae: Asopinae) is a predaceous stink bug. Life cycle of *A. malabarica* was studied in rice fields of east Vidarbha, India during Rabi season of 2018-20 on its prey, *Riptortus linearis* (Fabricius) (Heteroptera: Alydidae). The female laid 180 eggs in 4 to 8 batches on upper surface of leaves and each batch consisting of 20 to 45 eggs. The egg was barrel shaped, convex ventrally, flattened dorsally and measured 1.0 ± 0.047 mm, 0.76 ± 0.02 mm in length and width, respectively. The percentage of hatchability was very low (60%) and incubation period was 8.3 ± 0.33 days at 30 to 35 C. Pre-oviposition and oviposition period was 1 to 3 days, respectively. First, second, third, fourth, and fifth instar nymphal period were 4 ± 0.57 , 4.3 ± 0.33 , 5.3 ± 0.33 , 4.6 ± 0.33 and 5.6 ± 0.33 days, respectively. The mean consumptions of prey by second, third, fourth and fifth instar were 4.3 ± 0.33 , 8.6 ± 0.33 , 17.6 ± 1.4 and 26 ± 1.1 preys/bug, respectively, while adult fed on 39 ± 0.5 prey/bugs. In entire life a single *A. malabarica* killed 96.0 ± 3.7 preys. The nymphs of *A. malabarica* show cannibalism behavior in scarcity of food only. Total developmental period from egg to adult was 32.3 ± 2.02 days. Longevity of female was longer (11 to 27 days) than male (10-25 days). Present study provides important life history information of *A. malabarica* which is natural bio-agent against pests found in rice fields.

Keywords: *Amyotea malabarica*, Biology, Predatory efficiency, *Riptortus linearis*, Rice fields

Citation: Barsagade D.D. and Salwe A.A.: Biology of a pentatomid bug *Amyotea malabarica* (Fabricius) and its predatory efficiency in rice field of Vidarbha, India. Intern. J. Zool. Invest. 7 (Special Issue): 21-25, 2021.

<https://doi.org/10.33745/ijzi.2021.v07i02s.004>

Introduction

Amyotea malabarica (Fabricius) (Heteroptera: Pentatomidae: Asopinae) is a brilliantly reddish-coloured bug with black spots on dorsal surface and alternate white and black transverse lines on ventral side of abdomen. The bug is widely distributed in Bangladesh, Borneo, Burma, Celebes, China, India, Indonesia, Japan, Java, New Guinea, Phillipines, Sumatra, Sri Lanka and Taiwan in rice, cotton, soybean and other crop ecosystems

(Thomas, 1994). In India, it is recorded in several states (Salini and Viraktamath, 2015). Both nymphs and adults are predatory feeding by sucking the body juice of several insects, including rice pests such as rice bug *Leptocoris acuta* (Thunberg), *Nezara viridula* (Linnaeus) (Singh *et al.*, 1973) and lepidopteran larvae (Pati and Mathur, 1986). In Vidarbha (Maharashtra) *A. malabarica* was observed to consume a number of

परिवर्तना चा
वाटसरू

वर्ष २१ / अंक १५
किं. २५ रु. / १ ते १५ डिसेंबर २०२१

ISSN: 2250-3145

वर्गणीचे दर : वार्षिक : ६०० रु., त्रैवार्षिक : १,६०० रु.,
दशवार्षिक : ५,००० रु., सॉफ्ट कॉपीसाठी वार्षिक २०० रु.,
सॉफ्ट + छापील ७०० रु. वरील पत्त्यावर ई मनी ऑर्डर, चेक
किंवा डिमांड ड्राफ्टने वर्गणी पाठवावी. 'परिवर्तनाचा वाटसरू' या
नावाने चेक / डिमांड ड्राफ्ट असावा.

बँक ट्रान्सफरने वर्गणी भरण्यासाठी तपशील
बँक ऑफ इंडिया, फर्युसन कॉलेज रोड शाखा, पुणे । बचत खाते
: Pariwartanacha Vatsaru
बचत खाते क्र. : 051410110013969 । आयएफएससी क्र.
: BKID0000514 । एमआयसीआर क्र. : 411013014
Google Pay साठीचा मोबाईल नंबर : 8888708963
युपीआय पेमेंटसाठी बार कोड



युजीसी केअर ग्रुप १मध्ये 'परिवर्तनाचा वाटसरू' पाक्षिकाचा
मल्टिडिसिप्लिनरी या कोटीक्रमांतर्गत समावेश केला आहे.

प्रसाद वोंगळ चिवडिती।
दंभाचे आवडी बहिराट अंधळे।
सेवटासिं काळे होईल तोंड।।
सोन्यासेजारीं तो लाखेची जतन।
संतत ते गुण जैसे तैसे।।
सेव्यसेवकता न पडतां ठावी।
तुका म्हणे गोवी पावती हर्दी।।

- संत तुकाराम (अभंग क्र. १३९७)

सात्त्वाबादप्रमाणे १४व्या अखिल भारतीय मराठी साहित्य संमेलनाची चर्चा प्रसारमाध्यमांमधून जोरकसपणे झऱ्ह लागली आहे. नुकताच ब्राह्मण महासंघाने या साहित्य संमेलनात गीतकार जावेद अख्तर व गुलजार यांना निमंत्रण देण्यास कडाझून विरोध

लेखक तुमसर, जि. भंडारा येथे एस. एन. मोर कला-वाणिज्य मंहाविद्यालयात
मराठीचे सहाय्यक प्राध्यापक आहेत.
ईमेल- renukadas1983@gmail.com